

TITLE: Preparation of piperidine derivatives as tachykinin receptor antagonists for treatment of frequent urination and urinary incontinence

INVENTOR(S): Ikeura, Yoshinori; Hashimoto, Tadatoshi; Tarui, Naoki; Shirai, Junya; Yamashita, Masayuki

PATENT ASSIGNEE(S): Takeda Chemical Industries, Ltd., Japan

SOURCE: PCT Int. Appl., 264 pp.  
CODEN: PIXXD2


DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003101964	A1	20031211	WO 2003-JP6754	20030529
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2487688	A1	20031211	CA 2003-2487688	20030529
AU 2003241903	A1	20031219	AU 2003-241903	20030529
BR 2003011425	A	20050315	BR 2003-11425	20030529
EP 1553084	A1	20050713	EP 2003-733151	20030529
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
CN 1671662	A	20050921	CN 2003-818354	20030529
NZ 537330	A	20070427	NZ 2003-537330	20030529
JP 2004285038	A	20041014	JP 2003-154345	20030530
MX 2004PA11730	A	20050714	MX 2004-PA11730	20041125
US 20060167052	A1	20060727	US 2004-516252	20041129
ZA 2004010085	A	20060726	ZA 2004-10085	20041214
IN 2004KN01942	A	20061201	IN 2004-KN1942	20041216
NO 2004005701	A	20050216	NO 2004-5701	20041229
PRIORITY APPLN. INFO.:			JP 2002-159338	A 20020531
			JP 2003-17885	A 20030127
			WO 2003-JP6754	W 20030529
OTHER SOURCE(S):			MARPAT 140:27765	
GI				



AB The title compds. I [wherein Ar = (un)substituted aryl, aralkyl, or heteroaryl; R1 = H, acyl, (un)substituted hydrocarbyl, or heterocyclyl; X = O or (un)substituted NH; Z = (un)substituted CH2; ring A = (un)substituted piperidine; ring B = (un)substituted aryl; with exclusions] or prodrugs or salts thereof are prepared I have excellent tachykinin receptor antagonistic activity, and are useful for the treatment of frequent urination and urinary incontinence (no data). For example, the compound II•xHCl was prepared in a multi-step synthesis. II showed antagonistic activity with IC50 of 0.025 nM against human substance P receptor. Formulations containing I as an active ingredient were also described.

IT 632352-46-6P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of piperidine derivs. as tachykinin receptor antagonists for treatment of frequent urination and urinary incontinence)

RN 632352-46-6 CAPLUS

CN 3H-1,2,4-Triazol-3-one, 1,2-dihydro-5-[2-[(3R,4S)-4-[[[2-methoxy-5-[5-(trifluoromethyl)-1H-tetrazol-1-yl]phenyl]methyl]amino]-3-phenyl-1-piperidiny]-2-oxoethyl]-, rel- (CA INDEX NAME)

Relative stereochemistry.

